

RP01899

Leader in Biomolecular Solutions for Life Science



# Recombinant Human IFN-alpha I/IFN17 Protein

Catalog No.: RP01899 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	3451	P01571

### Tags

C-His

### Synonyms

Interferon alpha-17; IFN-alpha-17;  
Interferon alpha-88; Interferon alpha-I;  
LeIF I; Interferon alpha-T , IFNA17

## Product Information

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE.

### Endotoxin

< 0.01 EU/μg of the protein by LAL method

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Background

Interferons (IFN) are a family of cytokines with potent antiviral, antiproliferative and immunomodulatory properties, classified based on their binding specificity to cell surface receptors . Human IFNA2 was originally cloned in the early '80s and now more than a dozen closely related IFN alpha subtypes have been identified in both the human and mouse genome, each sharing about 80% amino acid (aa) sequence homology . Structurally, type I IFNs belong to the class of five helical bundle cytokines, with the IFNA subtypes containing 2 conserved disulfide bonds . There is not a mouse homolog for IFNA17, but mature human IFNA17 shares 58% aa sequence identity with chimpanzee IFNA17. The type I IFNs bind to the interferon alpha receptor (IFNAR), which consists of two subunits: IFNAR1 (alpha -subunit) and IFNAR2 (beta -subunit) . Individual IFNA subtypes are known to display unique efficacies to viral protection . IFNA17 has been shown to be potent against HIV-1 activity . Human IFNA17 is the only IFNA subtype identified with antiviral activity but a reduced ability to activate NK cells . A mutation in IFNA17, Ile184Arg, is associated with an increased risk for cervical cancer.

## Basic Information

### Description

Recombinant Human IFN-alpha I/IFN17 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Cys24-Asp189) of Human IFN-alpha I/IFN17 (Accession #NP\_067091.1) fused with a His tag at the C-terminus.

### Bio-Activity

Measured in a cell cytotoxicity assay using TF-1 cells. The ED<sub>50</sub> for this effect is 0.17 ± 0.68 ng/mL, corresponding to a specific activity of 1.47 × 10<sup>5</sup> - 5.88 × 10<sup>6</sup> units/mg.

### Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

### Operational Notes


For your safety and health, please wear a lab coat and disposable gloves for handling.

### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

\* For your safety and health, please wear a lab coat and disposable gloves when handling.

## Validation Data



Recombinant Human IFN-alpha I/IFN17 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Human IFN-alpha I/IFN17 Protein cytotoxicity assay using TF-1 cells. The ED<sub>50</sub> for this effect is 0.17 to 0.68 ng/mL, corresponding to a specific activity of  $1.47 \times 10^6 \sim 5.88 \times 10^5$  units/mg.