

Human Pancreatic CAFs – Stellate Cell

#Cat: NB-26-01967 Size: 1,000,000Cells

Catalog #: NB-26-01967

Cell #: >1x10⁶ cells

Storage: Liquid Nitrogen until ready for culture.

Product Format: Frozen Vial

While Culturing keep in 37°C CO₂ incubator

General Information

Carcinoma associated fibroblasts (CAFs) have recently been implicated in important aspects of epithelial solid tumor biology such as neoplastic progression, tumor growth, angiogenesis, and metastasis.

Product is for Research use only.

Frozen Vials are shipped in a Dry Ice Package.

Handling of Arriving Cells

1. Check all containers for leakage or breakage.
2. Remove the frozen cells from the dry ice packaging and immediately place the cells at a temperature below 130°C, preferably in liquid nitrogen vapor, until ready for use.
3. To ensure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

Medium

We recommend customers use our CAF Growth Media (cat. NB-26-02018) to culture these cells.

Protocol For Thawing the Cells

To ensure the highest level of viability, thaw the vial and initiate the culture as soon as possible upon receipt. If upon arrival, continued storage of the frozen culture is necessary, it should be stored in liquid nitrogen vapor phase and not at -70°C. Storage at -70°C will result in loss of viability.

1. Thaw the vial by gentle agitation in a 37°C water bath. To reduce the possibility of contamination, keep the O-ring and cap out of the water. Thawing should be rapid (approximately 2 minutes).
2. Remove the vial from the water bath as soon as the contents are thawed. Decontaminate by dipping in or spraying with 70% ethanol. All the operations from this point on should be carried out under strict aseptic conditions.
3. It is recommended that the cryoprotective agent be removed immediately. Centrifuge the cell suspension at approximately 125 x g for 5 to 10 minutes. Discard the supernatant and resuspend the cell pellet in an appropriate amount of fresh growth medium.
4. Add 6.0 to 8.0 mL of AlphaBioCoat (cat. NB-26-01690-20) to the T-Flask for 15 minutes. Aspirate the solution after 15 minutes, rinse with 8ml of 1XPBS. Discard the 1XPBS. Transfer the cells to an appropriate size T-Flask.

For Research Use Only

Neo Biotech Reagents Are For In Vitro And Certain Non-Human In Vivo Experimental Use Only and Not Intended for Use in Any Human Clinical Investigation, Diagnosis, Prognosis, Or Treatment. The Above Analyses Are Merely Typical Guides. They Are Not to Be Construed as Being Specifications. All Of the Above Information Is, To the Best of Our Knowledge, True and Accurate. However, Since the Conditions of Use Are Beyond Our Control, All Recommendations or Suggestions Are Made Without Guarantee, Express or Implied, On Our Part. We Disclaim All Liability in Connection with The Use of The Information Contained Herein or Otherwise, And All Such Risks Are Assumed by The User. We Further Expressly Disclaim All Warranties of Merchantability and Fitness for A Particular Purpose.

16/04/2025

Neo Biotech
74 rue des Suisses – 92000 Nanterre

5. It is important to avoid excessive alkalinity of the medium during recovery of the cells. It is suggested that, prior to the addition of the vial contents, the culture vessel containing the growth medium be placed into the incubator for at least 15 minutes to allow the medium to reach its normal pH (7.0 to 7.6).
6. Incubate the culture at 37°C in a suitable incubator. A 5% CO₂ in air atmosphere is recommended if using the medium described on this product sheet.

Subculturing Procedure

Note: If you have any questions or need clarification regarding the protocol for culturing these cells, please reach out to our technical department or call us at +33 9 77 40 09 09 with your questions before beginning.

Note: Volumes are given for a 75 cm² flask. Increase or decrease the amount of medium needed proportionally for culture vessels of other sizes.

1. Remove and discard culture medium.
2. Briefly rinse the cell layer with 1x PBS, remove and discard 1x PBS.
3. Add 2.0 to 3.0 mL of Cell Detachment solution (cat. NB-26-01693) to flask and observe cells under an inverted microscope until cell layer is dispersed (usually within 5 to 15 minutes).
 - **Note:** To avoid clumping do not agitate the cells by hitting or shaking the flask while waiting for the cells to detach. Cells that are difficult to detach may be placed at 37°C to facilitate dispersal.
4. Add 6.0 to 8.0 mL of AlphaBioCoat to the T-Flask for 15 minutes. Aspirate the solution after 15 minutes, rinse with 8ml of 1XPBS. Discard the 1XPBS.
5. Add 6.0 to 8.0 mL of complete growth medium and aspirate cells by gently pipetting.
6. To remove cell detachment solution, transfer cell suspension to centrifuge tube and spin at approximately 125 x g for 5 to 10 minutes. Discard supernatant and resuspend cells in fresh serum-free growth medium. Add appropriate aliquots of cell suspension to new culture vessels.
7. Place culture vessels in incubators at 37°C.

Subcultivation Ratio: A subcultivation ratio of 1:2 to 1:6 is recommended

Medium Renewal: Every 2 to 3 days



For Research Use Only

Neo Biotech Reagents Are For In Vitro And Certain Non-Human In Vivo Experimental Use Only and Not Intended for Use in Any Human Clinical Investigation, Diagnosis, Prognosis, Or Treatment. The Above Analyses Are Merely Typical Guides. They Are Not to Be Construed as Being Specifications. All Of the Above Information Is, To the Best of Our Knowledge, True and Accurate. However, Since the Conditions of Use Are Beyond Our Control, All Recommendations or Suggestions Are Made Without Guarantee, Express or Implied, On Our Part. We Disclaim All Liability in Connection with The Use of The Information Contained Herein or Otherwise, And All Such Risks Are Assumed by The User. We Further Expressly Disclaim All Warranties of Merchantability and Fitness for A Particular Purpose.

16/04/2025

Neo Biotech
74 rue des Suisses – 92000 Nanterre