

AS008

Leader in Biomolecular Solutions for Life Science



www.abclonal.com

# Cy3-conjugated Goat anti-Mouse IgG (H+L)

Catalog No.: AS008

67 Publications

## Basic Information

### Observed MW

### Calculated MW

### Category

Secondary Antibody

### Applications

IF/ICC,FC

### Cross-Reactivity

Mouse

### Conjugate

Cy3. Ex:548nm. Em:562nm.

## Background

Secondary antibodies are affinity-purified antibodies which will work with target-specific primary antibody in the detection, sorting or purification of its specified target. Secondary antibodies offer increased versatility enabling users to use many detection systems (e.g. HRP, AP, fluorescence). They can also provide greater sensitivity through signal amplification as multiple secondary antibodies. Most commonly, secondary antibodies are generated by immunizing the host animal (different from host species of primary antibody) with a pooled population of normal immunoglobulins from the host species of primary antibody and can be further purified and modified (i.e. antibody fragmentation, label conjugation, etc.) to ensure well-characterized specificity to corresponding normal immunoglobulins.

## Recommended Dilutions

IF/ICC 1:50 - 1:200

FC 1:100 - 1:800

## Immunogen Information

Gene ID

Swiss Prot

### Immunogen

This information is considered to be commercially sensitive.

### Synonyms

## Contact



[www.abclonal.com](http://www.abclonal.com)

## Product Information

### Source

Goat

### Isotype

Cy3 conjugated IgG

### Purification


Affinity purification

### Storage


Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS with 0.025% Sodium Azide,0.75% BSA,50% glycerol,pH7.3.


## Validation Data




Immunofluorescence analysis of HeLa cells using GAPDH Mouse mAb (AC033, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Mouse IgG (H+L)(AS008, dilution 1:200) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x.



Immunofluorescence analysis of NIH/3T3 cells using GAPDH Mouse mAb (AC033, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Mouse IgG (H+L)(AS008, dilution 1:200) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x.



Immunofluorescence analysis of PC-12 cells using GAPDH Mouse mAb (AC033, dilution 1:100) followed by a further incubation with Cy3 Goat Anti-Mouse IgG (H+L)(AS008, dilution 1:200) (Red). DAPI was used for nuclear staining (Blue). Objective: 40x.



Flow cytometry:  $1 \times 10^6$  K-562 cells (negative control, left) and Hep G2 cells (right) were surface-stained with Mouse Anti-Human GPC3 mAb ( $4 \mu\text{g}/\text{mL}$ , orange line) or secondary antibody only (blue line). Non-fluorescently stained HepG2 and K-562 cells were used as blank control (red line). Cy3 Goat Anti-Mouse IgG (H+L) (AS008, 1:200) was used as a secondary antibody.