

RP01322

Leader in Biomolecular Solutions for Life Science



Recombinant Rat TNF-alpha Protein

Catalog No.: RP01322

Recombinant

1 Publications

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	24835	P16599

Tags

C-His

Synonyms

DIF;TNF-alpha;TNFA;TNFSF2;cachexin;cachectin;TNF α ;TNFA

Product Information

Source	Purification
HEK293 cells	$\geq 95\%$ as determined by SDS-PAGE.

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μ m filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Tumor necrosis factor alpha is a cytokine produced primarily by monocytes and macrophages. It is found in synovial cells and macrophages in the tissues. The primary role of TNF α is in the regulation of immune cells. TNF α is able to induce apoptotic cell death, to induce inflammation, and to inhibit tumorigenesis and viral replication. Dysregulation of TNF α production has been implicated in a variety of human diseases, including major depression, Alzheimer's disease and cancer. Recombinant TNF α is used as an immunostimulant under the INN tasonermin. TNF-alpha is involved in fighting against the tumorigenesis, thus, is regarded as a molecular insight in cancer treatment.

Basic Information

Description

Recombinant Rat TNF-alpha Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Leu80-Leu235) of Rat TNFA (Accession #NP_036807.1) fused with an 6 \times His tag at the C-terminus.

Bio-Activity

Recombinant Rat TNF-alpha induces cytotoxicity in the L-929 Mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The ED₅₀ for this effect is typically 3.37-13.48 pg/mL, corresponding to a specific activity of 7.42 $\times 10^7$ ~2.97 $\times 10^8$ units/mg.

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Operational Notes


For your safety and health, please wear a lab coat and disposable gloves for handling.

Storage


Store at -20 $^{\circ}$ C. Store the lyophilized protein at -20 $^{\circ}$ C to -80 $^{\circ}$ C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week. Avoid repeated freeze/thaw cycles.

* For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



Recombinant Rat TNF-alpha Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Rat TNF-alpha induces cytotoxicity in the L-929 Mouse fibroblast cells in the presence of the metabolic inhibitor actinomycin D. The ED_{50} for this effect is typically 3.37-13.48 pg/mL, corresponding to a specific activity of $7.42 \times 10^7 \sim 2.97 \times 10^8$ units/mg.